arXiv:nucl-th/0611032v1 9 Nov 2006

COMPTON SCATTERING ON HE-3

D. CHOUDHURY* and D. R. PHILLIPS†

Department of Physics and Astronomy, Ohio University, Athens, Ohio 45701, USA
*E-mail: *choudhur@phy.ohiou.edu; †phillips@phy.ohiou.edu*

A. NOGGA

Institut für Kernphysik, Forschungszentrum Jülich, Jülich, Germany
E-mail: a.nogga@fz-juelich.de

We present first calculations for Compton scattering on ^{3}He. The objective of the calculation is an extraction of the neutron polarizabilities.

1. Introduction

Our goal is to devise ways to extract the neutron polarizabilities. Direct experiments on the neutron are not possible due to the lack of free neutron targets and this encouraged physicists to look at other avenues to extract information about the neutron polarizabilities. Elastic Compton scattering on ^{3}He is one such avenue and here we report on the first calculations of this process. Our results indicate that $\gamma\,^3$He scattering is indeed a promising way to extract the neutron polarizabilities.

2. The Calculation and Results

The irreducible amplitudes for the elastic scattering of real photons from the NNN system are first ordered and calculated in Heavy Baryon Chiral Perturbation Theory (HBχPT) to $\mathcal{O}(Q^3)$– they are the same as those computed in Ref. 1. These amplitudes are then sandwiched between the nuclear wavefunctions to finally obtain the scattering amplitude–

$$\mathcal{M} = \langle \Psi_f | \hat{O} | \Psi_i \rangle. \tag{1}$$

Using these amplitudes we calculate the differential cross-section (dcs) or the double polarization observables, Δ_z or Δ_x.[2]

In Fig. 1 shows the differential cross-section in the center of mass (c.m.) frame at 80 MeV. The left panel corresponds to calculations at different orders– $\mathcal{O}(Q^3)$, impulse approximation (IA) which is actually $\mathcal{O}(Q^3)$ but does not include two-body currents, and $\mathcal{O}(Q^3)$; and the right panel to varying the value of β_n around its $\mathcal{O}(Q^3)$ predicted value. Fig. 2 shows the variation in Δ_z (at 120 MeV in the c.m. frame) when $\Delta\gamma_{1n}(\times 10^{-4}\mathrm{fm}^4)$ is varied and that in Δ_x when $\Delta\gamma_{4n}(\times 10^{-4}\mathrm{fm}^4)$ is varied.



Fig. 1. The differential cross-section vs. c.m. angle at 80 MeV. Left panel shows $\frac{d\sigma}{d\Omega}$ calculated to different orders. Right panel shows the sensitivity of $\frac{d\sigma}{d\Omega}$ to $\Delta\beta_n(\times 10^{-4}\mathrm{fm}^3)$.



Fig. 2. Figure showing the sensitivity of the double-polarization observable, Δ_z to $\Delta\gamma_{1n}(\times 10^{-4}\mathrm{fm}^4)$ and of Δ_x to $\Delta\gamma_{4n}(\times 10^{-4}\mathrm{fm}^4)$ at 120 MeV in the c.m. frame.

Both Figs. 1 and 2 suggest sizeable sensitivity to the neutron polarizabilities in the $\gamma\,^3$He dcs and the double polarization observables. For a more detailed description of the calculation and results please refer to Ref. 2.

References

1. S. R. Beane, M. Malheiro, D. R. Phillips, and U. van Kolck, Nucl. Phys. **A656**, 367 (1999).
2. D. Choudhury, D. R. Phillips, and A. Nogga, (in preparation).